

Sm-Nd AND Rb-Sr AGES FOR NORTHWEST AFRICA 2977, A YOUNG LUNAR GABBRO FROM THE PKT.

L. E. Nyquist¹, C.-Y. Shih², Y. D. Reese³, and A. J. Irving⁴.

¹NASA Johnson Space Center, Houston, TX 77058. E-mail: laurance.e.nyquist@nasa.gov. ²JE-23, ESCG Jacobs-Sverdrup, Houston, TX 77058. ³JE-23, ESCG/Muniz Eng., Houston, TX 77058.

⁴Dept. Earth&Space Sci. Univ. Washington, Seattle, WA 98195.

Introduction: Northwest Africa (NWA) 2977 is an olivine gabbro cumulate equivalent to one of the lithologies in lunar mare breccia NWA 773 [1,2,3]. The ³⁹Ar-⁴⁰Ar age is 2.77±0.04 Ga based on the last ~57% of the gas release [4], similar to results for NWA 773 [5]. A Sm-Nd age (T) of 2.865±0.031 Ga and $\epsilon_{Nd} = -7.84 \pm 0.22$ for the NWA 773 gabbro reported by [6] has been revised to T = 2.993±0.032 Ga, $\epsilon_{Nd} = -4.5 \pm 0.3$ [7].

¹⁴⁷Sm-¹⁴³Nd isochron for NWA 2977: Whole rock, pyroxene, olivine, plagioclase, whole rock leachate (~phosphate) and the combined leachates from the mineral separates yield a well defined Sm-Nd isochron for an age T = 3.10±0.05 Ga and $\epsilon_{Nd,CHUR} = -3.74 \pm 0.26$ [8], or $\epsilon_{Nd,HEDR} = -4.61 \pm 0.26$ [9].

⁸⁷Rb-⁸⁶Sr isochron: NWA 2977 contains only a modest amount of Rb and/or Sr contamination. The Sr-isotopic composition of the contaminant closely resembles that of seawater. The whole rock residue after leaching combined with leach residues for plagioclase and pyroxene define an isochron age of 3.29±0.11 Ga for initial ⁸⁷Sr/⁸⁶Sr = 0.70287±18. The olivine residue, with lower Sr abundance of ~1.5 ppm, is only slightly displaced from the isochron. The relatively small uncertainties of the Rb-Sr isochron parameters and near-concordancy with the Sm-Nd age indicate that both the Rb-Sr and the Sm-Nd ages are reliable.

Discussion: The somewhat older Rb-Sr and Sm-Nd ages of NWA 2977 compared to its ³⁹Ar-⁴⁰Ar age suggest some radiogenic ⁴⁰Ar* loss from the sample analyzed by [4]. Initial ϵ_{Nd} for NWA 2977, when combined with ϵ_{Nd} for KREEP-rich samples, defines a line that when extrapolated passes between the CHUR and HEDR values at 4.568 Ga ago. Both types of samples lie within error limits of Nd-isotopic evolution for ¹⁴⁷Sm/¹⁴⁴Nd ~0.177 characteristic of KREEP-rich sources. Also, $\epsilon^{142}Nd$ for NWA 2977 is the same within error limits as for KREEP melt rock 14310 [10]. The ^{142,143}Nd isotopic evolution can be modeled for: (a) 50 Ma with chondritic ¹⁴⁷Sm/¹⁴⁴Nd (μ_{CHUR}) = 0.1967, (b) 168 Ma with $\mu_{moon} = 0.205$, (c) 1250 Ma with source region $\mu_S = 0.173$ (present-day values). A two-stage model for Sr-isotopic evolution gives ⁸⁷Rb/⁸⁶Sr ~0.207 in the NWA 2977 source, compared to ~0.18 in the rock and ~0.14-0.17 in the sources of A14 and A15 KREEP basalts. The NWA 2977 source evidently contained highly evolved Trapped Instantaneous Residual Liquid [11]. The age, Sr-and Nd-isotopes, and trace element abundances suggest an origin in the PKT (Procellarum KREEP Terrane).

References: [1] Bunch T. E. et al. 2006. *LPS XXXVII*, #1375. [2] Jolliff B. L. et al. 2007. *LPS XXXVIII*, #1489. [3] Zeigler R. A et al. 2007. *LPS XXXVIII*, #2109. [4] Burgess R. et al. 2007. *LPS XXXVIII*, #1603. [5] Fernandes V. A. et al. 2003. *Meteoritics & Planet. Sci.* 38: 555-564. [6] Borg L. et al. 2004. *Nature* 432: 209-211. [7] Borg L. et al. 2009. *Geochim. Cosmochim. Acta*, in press. [8] Jacobsen S. B. and Wasserburg G. J. 1984. *Earth & Planet. Sci. Lett.* 67: 137-150. [9] Nyquist L. et al. 2006. *Geochim. Cosmochim. Acta*, 70: 5990-6015. [10] Tou-boul M. et al. 2009. *LPS XL*, #2269. [11] Snyder G. A. and Taylor L. 1993. *Proc. NIPR Symp. Antarctic Met.* 6: 246-267.